

Energy Distribution of a Charged Regular Black Hole

I. Radinschi*

Department of Physics, “Gh. Asachi” Technical University,
Iasi, 6600, Romania

Abstract

We calculate the energy distribution of a charged regular black hole by using the energy-momentum complexes of Einstein and Møller.

Keywords: energy

PACS numbers: 04.20. Dw, 04.70. Bw

1 INTRODUCTION

One of the most interesting and intricate problem of relativity is the energy-momentum localization. The different attempts at constructing an energy-momentum density don't lead to a generally accepted expression. However, there are various energy-momentum complexes including those of Einstein [1], Tolman [2], Landau and Lifshitz [3], Papapetrou [4], Bergmann [5], Weinberg [6] and Møller [7]. Cooperstock [8] gave his opinion that the energy and momentum are confined to the regions of non-vanishing energy-momentum tensor of the matter and all non-gravitational fields. Although, the energy-momentum complexes are coordinate dependent they can give a reasonable result if calculations are carried out in Cartesian coordinates. Some interesting results obtained recently lead to the conclusion that different energy-momentum complexes give the same energy distribution for a given space-time [9]-[16].

*iradinsc@phys.tuiasi.ro

We calculate the energy distribution of a charged regular black hole by using the energy-momentum complexes of Einstein and Møller. We use the geometrized units ($G = 1, c = 1$) and follow the convention that Latin indices run from 0 to 3.

2 Energy in the Einstein prescription

The Reissner-Nordström (RN) metric is the only static and asymptotically flat solution of the Einstein-Maxwell equations and it represents an electrically charged black hole. The metric is given by

$$ds^2 = A(r)dt^2 - B(r)dr^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2) \quad (1)$$

where

$$A(r) = B^{-1}(r) = 1 - \frac{2M}{r} + \frac{q^2}{r^2} \quad (2)$$

and q and M are the electric charge and, respectively, the mass of the black hole.

A solution to the coupled system of the Einstein field and equations of the nonlinear electrodynamics was recently given by E. Ayón-Beato and A. Garcia (ABG) [17]. This solution represents a regular black hole with mass M and electric charge q and avoids thus the singularity problem. Also, the metric asymptotically behaves as the Reissner-Nordström solution. The usual singularity of the RN solution, at $r = 0$, has been smoothed out and now it simply corresponds to the origin of the spherical coordinates. The line element is given by (1) with

$$A(r) = B^{-1}(r) = 1 - \frac{2M}{r} \left(1 - \tanh\left(\frac{q^2}{2Mr}\right)\right). \quad (3)$$

If the electric charge vanishes we reach the Schwarzschild solution. At large distances (3) resembles to the Reissner-Nordström solution and can be written

$$A(r) = B^{-1}(r) = 1 - \frac{2M}{r} + \frac{q^2}{r^2} - \frac{q^6}{12M^2r^4} + O\left(\frac{1}{r^6}\right). \quad (4)$$

We obtain the energy distribution associated with the solution given by (1) and (3) in the Einstein and Møller prescriptions.

The Einstein energy-momentum complex [1] is given by

$$\Theta_i{}^k = \frac{1}{16\pi} H_i{}^{kl}, \quad (5)$$

where

$$H_i{}^{kl} = -H_i{}^{lk} = \frac{g_{in}}{\sqrt{-g}} [-g(g^{kn}g^{lm} - g^{ln}g^{km})]_{,l}. \quad (6)$$

$\Theta_0{}^0$ and $\Theta_\alpha{}^0$ are the energy and, respectively, the momentum components. The Einstein energy-momentum complex satisfies the local conservation laws

$$\frac{\partial \Theta_i{}^k}{\partial x^k} = 0. \quad (7)$$

The energy and momentum in the Einstein prescription are given by

$$P_i = \iiint \Theta_i{}^0 dx^1 dx^2 dx^3. \quad (8)$$

Using the Gauss theorem we obtain

$$P_i = \frac{1}{8\pi} \iint H_i{}^{0\alpha} n_\alpha dS, \quad (9)$$

where $n_\alpha = \left(\frac{x}{r}, \frac{y}{r}, \frac{z}{r}\right)$ are the components of a normal vector over an infinitesimal surface element $dS = r^2 \sin \theta d\theta d\varphi$.

The required nonvanishing components of $H_i{}^{kl}$ for the line element given by (1) and (3) are

$$\begin{aligned} H_0{}^{01} &= -\frac{4Mx(-1+\tanh(\frac{q^2}{2Mr}))}{r^3}, \\ H_0{}^{02} &= -\frac{4My(-1+\tanh(\frac{q^2}{2Mr}))}{r^3}, \\ H_0{}^{03} &= -\frac{4Mz(-1+\tanh(\frac{q^2}{2Mr}))}{r^3}. \end{aligned} \quad (10)$$

Using the above expressions in (9) we obtain

$$E(r) = M(1 - \tanh(\frac{q^2}{2Mr})). \quad (11)$$

From (11) it results that if $q = 0$ we have the energy of a Schwarzschild black hole.

After some calculations, we get the energy distribution of the ABG black hole

$$E(r) = M - \frac{q^2}{2r} + \frac{q^6}{24M^2r^3} - \frac{q^{10}}{240M^4r^5} + O\left(\frac{1}{r^6}\right). \quad (12)$$

Also, (12) can be written

$$E(r) = E_{RN}(r) + \frac{q^6}{24M^2r^3} - \frac{q^{10}}{240M^4r^5} + O\left(\frac{1}{r^6}\right). \quad (13)$$

3 Energy in the Møller prescription

The Møller energy-momentum complex [7] no needs to carry out calculations in Cartesian coordinates so we can calculate in any coordinate system.

Møller's energy-momentum complex is given by

$$M_i{}^k = \frac{1}{8\pi} \cdot \frac{\partial \chi_i{}^{kl}}{\partial x^l}, \quad (14)$$

where

$$\chi_i{}^{kl} = \sqrt{-g} \left(\frac{\partial g_{in}}{\partial x^m} - \frac{\partial g_{im}}{\partial x^n} \right) g^{km} g^{ln}. \quad (15)$$

The energy in the Møller prescription has the expression

$$E = \iiint M_0{}^0 dx^1 dx^2 dx^3 = \frac{1}{8\pi} \iiint \frac{\partial \chi_0{}^{0l}}{\partial x^l} dx^1 dx^2 dx^3. \quad (16)$$

For the line-element given by (1) and (3) the $\chi_0{}^{01}$ component is given by

$$\chi_0{}^{01} = \frac{(2Mr - 2Mr \tanh(\frac{q^2}{2Mr}) - q^2 + q^2 \tanh^2(\frac{q^2}{2Mr})) \sin \theta}{r}. \quad (17)$$

From (16) and (17) and applying the Gauss theorem we obtain the energy distribution

$$E(r) = M(1 - \tanh(\frac{q^2}{2Mr})) - \frac{q^2}{2r}(1 - \tanh^2(\frac{q^2}{2Mr})). \quad (18)$$

Also, from (18) we have

$$E(r) = M - \frac{q^2}{r} + \frac{q^6}{6M^2r^3} - \frac{q^{10}}{40M^4r^5} + O(\frac{1}{r^6}). \quad (19)$$

4 Discussion

Bondi [18] sustained that a nonlocalizable form of energy is not admissible in relativity so its location can in principle be found. We obtain the energy distribution of a charged regular black hole by using the energy-momentum complexes of Einstein and Møller. From (13) it results that in the Einstein prescription the first two terms in the expression of the energy correspond to the Penrose quasi-local mass definition (evaluated by Tod) [19]. The Møller formalism provides in the expression of the energy (19) a term $M - \frac{q^2}{r}$ which agrees with the Komar [20] prescription. Also, the Møller energy-momentum complex allows to make the calculations in any particular system of coordinates. However, the energy of the ABG black hole obtained by using the Einstein energy-momentum complex contains the quantity $M - \frac{q^2}{2r}$ which is agreeing with linear theory. Our result sustains the Virbhadra opinion [21] that "the Einstein energy-momentum complex is in a better bill of health" and gives acceptable results for many space-times. Also, Chang, Nester and Chen [22] showed that the energy-momentum complexes are actually quasilocal and legitimate expressions for the energy-momentum.

Acknowledgments

I am grateful to K. S. Virbhadra for his helpful suggestions.

References

- [1] C. Möller, *Ann. Phys. (NY)* **4**, 347(1958).
- [2] R. C. Tolman, *Relativity, Thermodynamics and Cosmology*, (Oxford Univ. Press, London 1934, p. 227).
- [3] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press, 1987, p. 280).
- [4] A. Papapetrou, *Proc. R. Irish. Acad.* **A52**, 11 (1948).
- [5] P. G. Bergmann and R. Thompson, *Phys. Rev.* **89**, 400-407 (1953)

- [6] S. Weinberg, Gravitation and Cosmology: Principles and Applications of General Theory of Relativity (John Wiley and Sons, Inc., New York, 1972, p. 165).
- [7] C. Möller, Ann. Phys. (NY) **4**, 347 (1958).
- [8] F. I. Cooperstock, Mod. Phys. Lett. **A14**, 1531 (1999).
- [9] K. S. Virbhadra, Phys. Rev. **D41**, 1086 (1990).
- [10] K. S. Virbhadra, Phys. Rev. **D42**, 2919 (1990).
- [11] N. Rosen and K. S. Virbhadra, Gen. Rel. Grav. **25**, 429 (1993).
- [12] K. S. Virbhadra, Pramana-J. Phys. **45**, 215 (1995).
- [13] A. Chamorro and K. S. Virbhadra, Pramana-J. Phys. **45**, 181 (1995).
- [14] J. M. Aguirregabiria, A. Chamorro and K. S. Virbhadra, Gen. Rel. Grav. **28**, 1393 (1996).
- [15] S. S. Xulu, Int. J. Mod. Phys. **D7**, 773 (1998), gr-qc/9910015
- [16] I-Ching Yang, Ching-Tzung Yeh, Rue-Ron Hsu and Chin-Rong Lee, Int. J. Mod. Phys. **D6**, (1997), 349.
- [17] E. Ayón-Beato and A. Garcia, Phys. Lett. **B464**, 25 (1999).
- [18] H. Bondi, Proc. R. Soc. London **A427**, 249 (1990).
- [19] R. Penrose, Proc. R. Soc. London **A381**, 53 (1982), K. P. Tod, Proc. R. Soc. London **A388**, 457 (1983).
- [20] A. Komar, Phys. Rev. **113**, 934 (1959).
- [21] K. S. Virbhadra, Phys. Rev. **D60**, 104041 (1999).
- [22] Chia-Chen Chang, J. M. Nester and Chiang-Mei Chen, Phys. Rev. Lett. **83**, 1897 (1999).